

AD-A164 691

ON THE EXISTENCE OF LI-YORKE POINTS IN THE THEORY OF  
CHAOS(U) ARMY BALLISTIC RESEARCH LAB ABERDEEN PROVING  
GROUND MD N P BHATIA ET AL. JAN 86 BRL-TR-2702


1/1

UNCLASSIFIED

F/G 12/1

NL





MICROCOPY RESOLUTION TEST CHART  
NATIONAL BUREAU OF STANDARDS-1963-A

AD-A164 691

AD

(2)

TECHNICAL REPORT BRL-TR-2702

ON THE EXISTENCE OF LI-YORKE  
POINTS IN THE THEORY OF CHAOS

Nam P. Bhatia  
Walter O. Egerland

January 1986

DTIC  
ELECTE  
FEB 24 1986  
S D

APPROVED FOR PUBLIC RELEASE; DISTRIBUTION UNLIMITED.

US ARMY BALLISTIC RESEARCH LABORATORY  
ABERDEEN PROVING GROUND, MARYLAND

DTIC FILE COPY

06 2 24 193

Destroy this report when it is no longer needed.  
Do not return it to the originator.

Additional copies of this report may be obtained  
from the National Technical Information Service,  
U. S. Department of Commerce, Springfield, Virginia  
22161.

The findings in this report are not to be construed as an official  
Department of the Army position, unless so designated by other  
authorized documents.

The use of trade names or manufacturers' names in this report  
does not constitute indorsement of any commercial product.

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER TECHNICAL REPORT BRL-TR-2702	2. GOVT ACCESSION NO. ADA 164691	3. RECIPIENT'S CATALOG NUMBER
4. TITLE (and Subtitle) On the Existence of Li-Yorke Points in the Theory of Chaos.		5. TYPE OF REPORT & PERIOD COVERED
		6. PERFORMING ORG. REPORT NUMBER
7. AUTHOR(s) Nam P. Bhatia Walter O. Egerland		8. CONTRACT OR GRANT NUMBER(s)
9. PERFORMING ORGANIZATION NAME AND ADDRESS US Army Ballistic Research Laboratory ATTN: SLCBR-SE Aberdeen Proving Ground, MD 21005-5066		10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS  RDT&E 1L162618AH43
11. CONTROLLING OFFICE NAME AND ADDRESS US Army Ballistic Research Laboratory ATTN: SLCBR-DD-T Aberdeen Proving Ground, MD 21005-5066		12. REPORT DATE January 1986
		13. NUMBER OF PAGES 12
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office)		15. SECURITY CLASS. (of this report)  UNCLASSIFIED
		15a. DECLASSIFICATION/DOWNGRADING SCHEDULE
16. DISTRIBUTION STATEMENT (of this Report)  Approved for public release; distribution unlimited.		
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)		
18. SUPPLEMENTARY NOTES The first author is Professor of Mathematics at the University of Maryland, Baltimore County. The results were established under the U.S. Army Summer Faculty Research and Engineering Program and will be published in the Journal of Nonlinear Analysis, Pergamon Press LTD., Oxford, England.		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number) Theory of Chaos Li-Yorke Inequalities and Points Non-periodic Conditions for Chaos		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number) The report establishes an addendum to the Li-Yorke theorem, lists the complete set of Li-Yorke inequalities, and provides in the Equivalence theorem non-periodic conditions equivalent to the known periodic conditions that guarantee chaos. <i>continuous functions</i>		

# TABLE OF CONTENTS

	Page
I. INTRODUCTION.....	5
II. DEFINITIONS AND NOTATIONS.....	5
III. AN EXAMPLE.....	6
IV. THE RELATIONSHIP BETWEEN LI-YORKE POINTS AND POINTS OF PERIOD THREE.....	6
V. REFERENCES .....	9
VI. DISTRIBUTION LIST .....	11

Accession For	
NTIS CRA&I	<input checked="" type="checkbox"/>
DTIC TAB	<input type="checkbox"/>
Unannounced	<input type="checkbox"/>
Justification .....	
By .....	
Distribution /	
Availability Codes	
Dist	Avail and/or Special
A-1	

## I. INTRODUCTION

Li and Yorke<sup>1</sup> introduced in their fundamental paper the term "chaotic" for a class of self-mappings of an interval. The real function  $f$  is chaotic if (a) there are points of arbitrarily large periods and (b) there is an uncountable set  $S$  such that for every  $x_0, y_0 \in S$ ,  $x_0 \neq y_0$ ,  $\limsup_{n \rightarrow \infty} |f^n(x_0) - f^n(y_0)| > 0$  and  $\liminf_{n \rightarrow \infty} |f^n(x_0) - f^n(y_0)| = 0$ . Following the Li-Yorke result that "period three implies chaos," many authors worked on periodic conditions that allow the same conclusion. The best known of these conditions is "period  $p \neq 2^a$  implies chaos." Such investigations are summarized in Targonski's monograph.<sup>2</sup>

Li and Yorke also introduced four-point inequalities satisfied by a point and its three successors with respect to the given function  $f$ . They showed that these imply the existence of a three-period and hence chaos. Our investigations in this paper show that the Li-Yorke inequalities play a fundamental role in the theory of chaos. We, therefore, formalize the notion of a Li-Yorke point and obtain non-periodic conditions equivalent to the known periodic conditions that guarantee chaos. The importance of the results resides in the fact that the set of Li-Yorke points is always an open set, whereas the number of periodic points of a given period  $p \neq 2^a$  is in general countable.

## II. DEFINITIONS AND NOTATIONS

Let  $f: R \rightarrow R$  be continuous. If  $x_0 \in R$ , the orbit of  $x_0$  under  $f$  is defined as the set  $\{x: x = f^n(x_0), n = 0, 1, \dots\}$ , where, for every positive integer  $n$ ,  $f^n$  is the  $n$ -th iterate of  $f$  and  $f^0(x_0) = x_0$ . We shall write  $x_n := f^n(x_0)$  for a given  $x_0 \in R$  and call  $x_1, x_2, \dots$  the successors of  $x_0$ . A pre-orbit of a given  $x_0 \in R$  is any (finite or infinite) sequence  $x_0, x_{-1}, x_{-2}, \dots$  such that  $f(x_{-n}) = x_{-(n-1)}$  for all  $n$  for which  $x_{-n}$  is defined. The points  $x_{-1}, x_{-2}, \dots$  in any such sequence are called predecessors of  $x_0$ . A point  $x_0$  is called critical if  $f(x_0) = x_0$ , i.e., a critical point of  $f$  is a fixed point of  $f$ . A periodic point  $x_0$  of period  $p > 1$  ( $p$  a positive integer) is a point for which the relations  $f^p(x_0) = x_0$ ,  $f^k(x_0) \neq x_0$ ,  $1 \leq k < p$ , hold.

The following fundamental results are now well-known.

**Theorem** (Sarkovskii).<sup>3</sup> For  $m, n = 0, 1, \dots$  consider the total ordering of the positive integers:

$$3 < 5 < 7 < \dots < 2 \cdot 3 < 2 \cdot 5 < 2 \cdot 7 \dots < 2^n \cdot 3 < 2^n \cdot 5 < 2^n \cdot 7 \\ < \dots < 2^m < 2^{m-1} < \dots < 2^2 < 2 < 1.$$

If a continuous mapping  $f: R \rightarrow R$  has a periodic point of period  $p$ , then it also has a periodic point of period  $q$  for every  $q > p$  (in the above total order).

**Theorem** (Li-Yorke): Let  $f: R \rightarrow R$  be continuous. If there is a point  $x_0 \in R$  such that either  $x_3 < x_0 < x_1 < x_2$  or  $x_3 > x_0 > x_1 > x_2$ , then  $f$  has a point of period three.

<sup>1</sup> T.-Y. Li and J. A. Yorke, "Period three implies Chaos," *Amer. Math. Monthly* 82 (1975), pp. 985-992.

<sup>2</sup> Gyorgy Targonski, "Topics in Iteration Theory," *Studia Mathematica. Skript 6*, Vandenhoek and Ruprecht, Gottingen and Zurich (1981).

<sup>3</sup> A. N. Sarkovskii, "Coexistence of Cycles of a Continuous Map of a Line into Itself," *Ukrain. Mat. Zh.* 16 (1964), pp. 61-71.

**Theorem (Li-Yorke):** Let  $f : R \rightarrow R$  be continuous. If there is a point  $x_0 \in R$  such that either  $x_3 < x_0 < x_1 < x_2$  or  $x_3 > x_0 > x_1 > x_2$ , then  $f$  has a point of period three. Furthermore, if  $f$  has a three periodic point, there exists an uncountable set  $S \subset R$  such that for every  $x_0, y_0 \in S$ ,  $x_0 \neq y_0$ ,

$$\limsup_{n \rightarrow \infty} |x_n - y_n| > 0$$

and

$$\liminf_{n \rightarrow \infty} |x_n - y_n| = 0.$$

**Definition.** A point  $x_0 \in R$  is called a Li-Yorke point of  $f : R \rightarrow R$  if  $x_0$  satisfies a Li-Yorke inequality

$$x_3 < x_0 < x_1 < x_2$$

or

$$x_3 > x_0 > x_1 > x_2.$$

### III. AN EXAMPLE

The following example shows that the existence of a three periodic point does not guarantee the existence of a Li-Yorke point. The quadratic mapping  $g(y) = ay^2 + 2by + c$ ,  $a \neq 0$ ,  $b, c$ , real constants, may be brought in the form  $f(x) = x^2 - r$  by setting  $y = a^{-1}(x - b)$ ,  $g = a^{-1}(f - b)$ , and  $r = b^2 - b - ac$ .  $f$  has a three periodic orbit for  $r = 7/4$ , but no point  $x_0 \in R$  is a Li-Yorke point for  $r \leq 7/4$ .

### IV. THE RELATIONSHIP BETWEEN LI-YORKE POINTS AND POINTS OF PERIOD THREE

We apply frequently the following crossing property of real continuous functions.

**Lemma.** Let  $f(x)$  be continuous on  $[a, b]$ . If either

$$f(a) \geq a, \quad f(b) \leq b$$

or

$$f(a) \leq a, \quad f(b) \geq b,$$

then there exists  $x \in [a, b]$  such that  $x = f(x)$ . In particular, the existence of such an  $x \in [a, b]$  follows from the inclusions  $[a, b] \subset [f(a), f(b)]$  or  $[a, b] \supset [f(a), f(b)]$ .

We shall denote the  $k$ -th iterate of  $x_0$  under the function  $f^m$  by  $x_k^m$ ,  $k = 0, 1, \dots$ . Thus  $x_k^m = (f^m)^k(x_0) = x_{mk}$  and, in particular,  $x_0^m = x_k^0 = x_0$  for all nonnegative integers  $k$  and  $m$ .

**Theorem 1.** If  $f$  has a three periodic point, then  $f^2$  has a Li-Yorke point.



**Proof.** Let  $y_0$  be a three periodic point of  $f$ . Thus  $y_0 = y_3$  and  $y_1 \neq y_k$ ,  $0 \leq i < k \leq 3$ . We can assume without loss of generality that  $y_0 < y_1 < y_2$ . Since  $f([y_1, y_2]) \supset [y_0, y_2] \supset [y_1, y_2]$ , there is a critical point  $c_0 \in (y_1, y_2)$ . Hence

$$y_0 < y_1 < c_0 < y_2.$$

Since  $f(y_0) = y_1 < c_0$  and  $f(y_1) = y_2 > c_0$ , we have a  $c_{-1} \in (y_0, y_1)$ , and since  $f(y_2) = y_0 < c_{-1}$  and  $f(c_0) = c_0 > c_{-1}$ , there is a  $c_{-2} \in (c_0, y_2)$ , thus yielding the inequality

$$y_0 < c_{-1} < y_1 < c_0 < c_{-2} < y_2.$$

Similar reasoning establishes points  $c_{-j}$ ,  $j = 3, 4, 5, 6$ , and the inequality

$$y_0 < c_{-1} < y_1 < c_{-3} < c_{-6} < c_0 < c_{-6} < c_{-4} < c_{-2} < y_2.$$

If we identify  $c_{-6}$  with  $x_0$ , we obtain  $c_{-6} = x_0$ ,  $c_{-4} = x_1^2$ ,  $c_{-2} = x_2^2$ ,  $c_0 = x_3^2$  and the inequality  $x_3^2 < x_0 < x_1^2 < x_2^2$ . This completes the proof.

**Theorem 2.** If  $f$  has a Li-Yorke point  $x_0$ ,  $f$  has at least two distinct three periodic orbits.

**Proof.** Let  $S = \{x_0 : x_3 < x_0 < x_1 < x_2 \text{ or } x_2 < x_1 < x_0 < x_3\}$ . Then  $S$  is non-empty by hypothesis and open because  $f$  is continuous. Each component of  $S$  is an open interval. If  $I = (a_0, b_0)$  is a component of  $S$ , we will show that  $-\infty < a_0 < b_0 < \infty$  and that both  $a_0$  and  $b_0$  are three periodic points in different orbits. Fix  $x_0 \in I$  and assume that  $x_3 < x_0 < x_1 < x_2$ . Then the same inequality holds for all  $x_0 \in I$ . Assuming first that  $-\infty < a_0 < b_0 < \infty$ , we note that  $a_0$  and  $b_0$  are limit points of points  $x_0$  for which  $x_3 < x_0 < x_1 < x_2$ . Hence by continuity of  $f$  we must have the inequalities  $a_3 \leq a_0 \leq a_1 \leq a_2$  and  $b_3 \leq b_0 \leq b_1 \leq b_2$  with at least one equality sign holding in each of them. We claim that  $a_3 = a_0 < a_1 < a_2$  and  $b_3 = b_0 < b_1 < b_2$ . Noting that  $f(x_1) = x_2 > x_1$  and  $f(x_2) = x_3 < x_2$ , it follows that there is a critical point  $c_0 \in (x_1, x_2)$ , so that  $x_3 < x_0 < x_1 < c_0 < x_2$ . Since  $f(x_0) = x_1 < c_0$  and  $f(x_1) = x_2 > c_0$  imply the existence of  $c_{-1} \in (x_0, x_1)$  with  $f(c_{-1}) = c_0$ , we have  $x_3 < x_0 < c_{-1} < x_1 < c_0 < x_2$ . As  $a_0 < x_0$  and  $f(a_0) = a_1 = a_0 < c_{-1}$ ,  $f(x_0) = x_1 > c_{-1}$ , we have  $c_{-2} \in (a_0, x_0) \subset I$  with  $f(c_{-2}) = c_{-1}$ . But  $c_{-2} < c_{-1} < c_0 = c_1$  yields by relabelling a  $y_0 = c_{-2} \in I$  such that  $y_0 < y_1 < y_2 = y_3$ , a contradiction. Hence  $a_0 < a_1$ . If now  $a_1 = a_2$ , we have  $a_0 < a_1 = a_2 = a_3$  which contradicts  $a_3 \leq a_0 \leq a_1 \leq a_2$ . Therefore, we must have  $a_1 < a_2$ , and hence  $a_3 = a_0 < a_1 < a_2$ , i.e.,  $a_0$  is a three periodic point. Similarly,  $b_0$  is a three periodic point for which  $b_3 = b_0 < b_1 < b_2$  holds. Since  $a_0 < b_0$ ,  $a_0$  is not in the orbit of  $b_0$ , and so the orbits of  $a_0$  and  $b_0$  are distinct. It remains to show that  $a_0 \neq -\infty$  and  $b_0 \neq \infty$ . Since  $x_0 < b_0$  and  $(x_0, b_0) \subset I$  and  $c_{-1} \notin S \supset I$ , we must have  $x_0 < b_0 < c_{-1}$  as  $x_0 < c_{-1}$ . This shows that  $b_0 < \infty$ . To see that  $a_0 > -\infty$ , we note that  $x_3 \in f^2[c_{-1}, c_0]$ , i.e.,  $x_3$  lies in a compact set. If  $A = \inf f^2[c_{-1}, c_0]$ , then for  $x_0 < A$  we conclude  $x_0 < x_3$  which contradicts  $x_0 \in I$  with  $a_0 = -\infty$ . This completes the proof.

The theorem extends the Li-Yorke theorem and may be illustrated by the example  $f(x) = x^2 - 2$ . The point  $x_0 = \sqrt{2}$  is a Li-Yorke point of  $f$  and  $(2 \cos \frac{2}{7}\pi, 2 \cos \frac{4}{7}\pi, 2 \cos \frac{8}{7}\pi)$ ,  $(2 \cos \frac{2}{9}\pi, 2 \cos \frac{4}{9}\pi, 2 \cos \frac{8}{9}\pi)$  are two distinct three periodic orbits.

The following theorem extends the Li-Yorke theorem in another direction. It establishes equivalent companion inequalities to the Li-Yorke inequalities.

**Theorem 3.** The sets of points  $A = \{x_0 : x_3 < x_0 < x_1 < x_2\}$ ,  $B = \{x_0 : x_1 < x_2 < x_0 < x_3\}$ , and  $C = \{x_0 : x_2 < x_3 < x_0 < x_1\}$  are either all empty or all non-empty. The same statement holds for the sets in which all inequalities are reversed.

**Proof.** Let  $A \neq \emptyset$ . Then for some  $x_0 \in R$  we have  $x_3 < x_0 < x_1 < x_2$ . Since  $f(x_1) = x_2 > x_0$  and  $f(x_2) = x_3 < x_0$ , there is  $x_{-1} \in (x_1, x_2)$  such that  $x_0 < x_1 < x_{-1} < x_2$ . Upon relabelling  $y_0 = x_{-1}$ , we have  $y_1 < y_2 < y_0 < y_3$ , so that  $B \neq \emptyset$ , i.e.,  $A \neq \emptyset$  implies  $B \neq \emptyset$ . Let  $B \neq \emptyset$ . Then, if  $x_1 < x_2 < x_0 < x_3$ , it follows from  $f(x_0) = x_1 < x_0$  and  $f(x_2) = x_3 > x_2$  that there is a critical point  $c_0 \in (x_2, x_0)$  such that  $x_1 < x_2 < c_0 < x_0 < x_3$ . Since  $f(c_0) = c_0 < x_0$  and  $f(x_2) = x_3 > x_0$ , there is  $x_{-1} \in (x_2, x_0)$ , so that  $x_1 < x_2 < x_{-1} < x_0$ . Letting  $y_0 = x_{-1}$ , we have  $y_2 < y_3 < y_0 < y_1$ . Thus  $B \neq \emptyset$  implies  $C \neq \emptyset$ . Finally, assume  $C \neq \emptyset$ . If for some  $x_0 \in R$   $x_2 < x_3 < x_0 < x_1$ , then there exists  $x_{-1} \in (x_2, x_0)$  since  $f(x_2) = x_3 < x_0$  and  $f(x_0) = x_1 > x_0$ . From  $x_2 < x_{-1} < x_0 < x_1$  follows, setting  $x_{-1} = y_0$ ,  $y_3 < y_0 < y_1 < y_2$ , and hence that  $C \neq \emptyset$  implies  $A \neq \emptyset$ . This completes the proof of the theorem.

Examples show that other four-point inequalities between  $x_0, x_1, x_2, x_3$  do not assure the existence of three-periodic orbits. Therefore, the theorem lists the complete set of Li-Yorke four-point inequalities.

**Theorem 4 (Equivalence Theorem).** The following statements are equivalent.

- (a)  $f$  has a point of period  $p \neq 2^n$ .
- (b) There exists  $m \geq 1$  such that  $f^m$  has a point of period three.
- (c)  $f^{2m}$  has Li-Yorke points.
- (d)  $f^{2m}$  has at least two three-periodic orbits.

**Proof.** If  $f$  has a point of period  $p \neq 2^n$ , then  $f$  has a point of period  $3 \cdot 2^k$  for some  $k \geq 0$  by the theorem of Sarkovskii. Such a point is a three-periodic point of  $f^m$ ,  $m = 2^k$ . Hence (a) implies (b). (b) implies (c) by Theorem 1, and (c) implies (d) by Theorem 2. Now let  $x_0$  be a three-periodic point of  $f^q$ ,  $q = 2m$ ,  $x_0 = f^{3q}(x_0)$ ,  $x_0 \neq f^q(x_0)$ ,  $x_0 \neq f^{2q}(x_0)$ .  $x_0$  is thus a periodic point of  $f$  of minimal period  $N$ ,  $2 \leq N \leq 3q$ . This implies that  $x_0 = f^{kN}(x_0)$  for every  $k \geq 1$  and  $x_0 \neq f^s(x_0)$  if  $s \neq kN$ ,  $k = 1, 2, \dots$ . Hence  $3q = k'N$  for some  $k' \geq 1$ . This shows that either 3 divides  $k'$  or 3 divides  $N$ . If 3 divides  $k'$ , then  $q = sN$  and consequently  $f^q(x_0) = x_0 = f^N(x_0)$ , a contradiction. Hence 3 divides  $N$ . But then  $N \neq 2^n$  for every  $n \geq 0$ . Therefore,  $f$  has period  $N \neq 2^n$  and (d) implies (a). This completes the proof of the Equivalence Theorem.

## V. REFERENCES

1. T.-Y. Li and J. A. Yorke, "Period three implies Chaos.," Amer. Math. Monthly 82 (1975), pp. 985-992.
2. Gyorgy Targonski, "Topics in Iteration Theory," Studia Mathematica, Skript 6, Vandenhoeck and Ruprecht, Gottingen and Zurich (1981).
3. A. N. Sarkovskii, "Coexistence of Cycles of a Continuous Map of a Line into Itself," Ukrain. Mat. Zh. 16 (1964), pp. 61-71.

# DISTRIBUTION LIST

<u>Copies</u>	<u>Organization</u>	<u>Copies</u>	<u>Organization</u>
12	Administrator Defense Technical Info Center ATTN: DTIC-DDA Cameron Station Alexandria, VA 22304-6145	1	Commander US Army Aviation Research and Development Command ATTN: AMSAV-E 4300 Goodfellow Blvd St. Louis, MO 63120
1	HQDA DAMA-ART-M Washington, DC 20310	1	Director US Army Air Mobility Research and Development Laboratory Ames Research Center Moffett Field, CA 94035
1	Commander US Army Materiel Command ATTN: AMCDRA-ST 5001 Eisenhower Avenue Alexandria, VA 22333-0001	1	Commander US Army Communications - Electronics Command Fort Monmouth, NJ 07703
10	Central Intelligence Agency Office of Central Reference Dissemination Branch Room GE-47 HOS Washington, D.C. 20502	1	Commander ERADCOM Technical Library ATTN: DELSD-L (Reports Section) Fort Monmouth, NJ 07703-5301
1	Commander Armament R&D Center US Army AMCCOM ATTN: SMCAR-TSS Dover, NJ 07801	1	Commander US Army Missile Command Research, Development, and Engineering Center ATTN: AMSMI-RD Redstone Arsenal, AL 35898
1	Commander Armament R&D Center US Army AMCCOM ATTN: SMCAR-TDC Dover, NJ 07801	1	Director US Army Missile and Space Intelligence Center ATTN: AIAMS-YDL Redstone Arsenal, AL 35898
1	Director Benet WEapons Laboratory Armament R&D Center US Army AMCCOM ATTN: SMCAR-LCB-TL Watervliet, NY 12189	1	Commander US Army Tank Automotive Command ATTN: AMSTA-TSL Warren, MI 48090
1	Commander US Army Armament, Munitions and Chemical Command ATTN: SMCAR-ESP-L Rock Island, IL 61299	1	Director US Army TRADOC Systems Analysis Activity ATTN: ATAA-SL White Sands Missile Range, NM 88002

## DISTRIBUTION LIST

<u>Copies</u>	<u>Organization</u>
1	Commandant US Army Infantry School ATTN: ATSH-CD-CSO-OR Fort Benning, GA 31905
1	Commander US Army Development and Employment Agency ATTN: MODE-TED-SAB Fort Lewis, WA 98433
1	AFWL/SUL Kirtland AFB, NM 87117
1	Air Force Armament Laboratory ATTN: AFATL/DLODL Eglin AFB, FL 32542-5000

### Aberdeen Proving Ground

Dir, USAMSAA  
ATTN: AMXSY-D  
AMXSY-MP, H. Cohen  
Cdr, USATECOM  
ATTN: AMSTE-TO-F  
Cdr, CRDC, AMCCOM  
ATTN: SMCCR-RSP-A  
SMCCR-MU  
SMCCR-SPS-IL

## USER EVALUATION SHEET/CHANGE OF ADDRESS

This Laboratory undertakes a continuing effort to improve the quality of the reports it publishes. Your comments/answers to the items/questions below will aid us in our efforts.

1. BRL Report Number \_\_\_\_\_ Date of Report \_\_\_\_\_

2. Date Report Received \_\_\_\_\_

3. Does this report satisfy a need? (Comment on purpose, related project, or other area of interest for which the report will be used.) \_\_\_\_\_  
\_\_\_\_\_  
\_\_\_\_\_

4. How specifically, is the report being used? (Information source, design data, procedure, source of ideas, etc.) \_\_\_\_\_  
\_\_\_\_\_  
\_\_\_\_\_

5. Has the information in this report led to any quantitative savings as far as man-hours or dollars saved, operating costs avoided or efficiencies achieved, etc? If so, please elaborate. \_\_\_\_\_  
\_\_\_\_\_  
\_\_\_\_\_

6. General Comments. What do you think should be changed to improve future reports? (Indicate changes to organization, technical content, format, etc.) \_\_\_\_\_  
\_\_\_\_\_  
\_\_\_\_\_

CURRENT ADDRESS	_____
	Name
	_____
	Organization
	_____
	Address
	_____
	City, State, Zip

7. If indicating a Change of Address or Address Correction, please provide the New or Correct Address in Block 6 above and the Old or Incorrect address below.

OLD ADDRESS	_____
	Name
	_____
	Organization
	_____
	Address
	_____
	City, State, Zip

(Remove this sheet along the perforation, fold as indicated, staple or tape closed, and mail.)

----- FOLD HERE -----

Director  
U.S. Army Ballistic Research Laboratory  
ATTN: SLCBR-DD-T  
Aberdeen Proving Ground, MD 21005-5066




NO POSTAGE  
NECESSARY  
IF MAILED  
IN THE  
UNITED STATES

OFFICIAL BUSINESS  
PENALTY FOR PRIVATE USE, \$300

**BUSINESS REPLY MAIL**  
FIRST CLASS PERMIT NO 12062 WASHINGTON, DC  
POSTAGE WILL BE PAID BY DEPARTMENT OF THE ARMY

Director  
U.S. Army Ballistic Research Laboratory  
ATTN: SLCBR-DD-T  
Aberdeen Proving Ground, MD 21005-9989



----- FOLD HERE -----

DTIC

END

4-86